arXiv:1105.5179v3 [math.AC] 13 Jun 2011

The structure of finite local principal ideal rings

Tongsuo Wu[a*], Houyi Yu[b†] and Dancheng Lu[c‡]
[a,b]Department of Mathematics, Shanghai Jiaotong University
Shanghai 200240, P. R. China
[c]Department of Mathematics, Suzhou University, Suzhou 215006, P.R. China

Abstract. A ring R is called a PIR, if each ideal of R is a principal ideal. A local ring $(R, \mathfrak{m})$ is an artinian PIR if and only if its maximal ideal $\mathfrak{m}$ is principal and has finite nilpotency index. In this paper, we determine the structure of a finite local PIR.

Key Words: finite local ring, polynomial rings, ideal structure, ring extension

1 Introduction

Throughout this paper, all rings R are assumed to be commutative with identity 1_R. For a ring R, let $U(R)$ be the set of invertible elements of R, $J(R)$ the Jacobson radical of R and $\mathrm{char}(R)$ the characteristic of R. A ring R is called a PIR if each ideal of R is a principal ideal. For a local ring $(R, \mathfrak{m})$ with a finite number of ideals (i.e., an artinian local ring R), R is a PIR if and only if the maximal ideal $\mathfrak{m}$ is cyclically generated ([1, Proposition 8.8] or [5, Theorem 2.1]). Furthermore, the number of nontrivial ideals of R is s if $s+1$ the nilpotency index of $\mathfrak{m}$. For any local ring $(R, \mathfrak{m}, \mathfrak{k})$ with $\mathrm{char}(\mathfrak{k}) = p$ and any pair $x, y \in \mathfrak{m}$, note that $(x, y) = (x + my, x + ny)$ holds for all integers n, m satisfying $0 \leq m < n \leq p - 1$, thus R has at least $p + 2$ nontrivial ideals if (x, y) is not cyclic: $(x, y), (y), (x + my)$ $(0 \leq m \leq p - 1)$. Now for any integer $r \geq 4$, let $\mathbb{I}(r)$ be the set of finite local rings with r nontrivial ideals. Then for any $(R, \mathfrak{m}, \mathfrak{k}) \in \mathbb{I}(r)$, if $\mathrm{char}(\mathfrak{k}) \geq r - 1$,

---

[*] Corresponding author, tswu@sjtu.edu.cn (T. Wu)
[†] yhy178@163.com (H.Y. Yu)
[‡] ludancheng@suda.edu.cn (D. Lu)

then R is a PIR. This implies that almost all finite local rings are PIR. In particular, a finite local ring $(R, \mathfrak{m}, \mathfrak{k})$ with four nontrivial ideals is always a PIR if $\mathrm{char}(\mathfrak{k}) \geq 3$.

Let p be a prime number. Recall from [5], page 255 (or [2], Lemma 13, page 79 and Corollary 2, page 83.) that if P is a field of characteristic p, then there exists an unramified complete discrete valuation ring, V, *of characteristic* 0 whose residue field is isomorphic to P. This ring V is uniquely determined up to isomorphism by P and is called the *v-ring with residue field* P. In [5, Theorem 3.2], the author uses the Cohen structure theory of complete local rings (see [2]) to prove that a local ring $(R, \mathfrak{m})$ is an artinian PIR, with $pR = \mathfrak{m}^m$, and with $\mathfrak{m}$ of nilpotency index s, if and only if $R \cong V[X]/(f(X), X^s)$ for some Eisenstein polynomial $f(X) \in V[X]$ of degree m, where V is any v-ring with residue field $R/\mathfrak{m}$. Moreover, in [5, Theorems 3.5, 3.6] the author classifies all tamely ramified (i.e., $p \nmid m$ in the aforementioned theorem) artinian local PIRs up to isomorphism by taking advantage of the established results on v-rings.

The purpose of this paper is to characterize a finite local PIR $(R, \mathfrak{m})$ as some homomorphic image of $\mathbb{Z}_{p^r}[X, Y]$ and we determine the kernel, where $\mathrm{char}(R) = p^r$. In Theorem 2.2, for any finite local ring R, we construct a coefficient field of R starting from a generator of the multiplicative cyclic group $(R/\mathfrak{m})^*$. The main structural results are Corollary 2.3 and Theorem 3.1. Note that our structure theorems as well as the proofs are quite different from [5].

2 Finite local rings with a prime characteristic

In this section, we classify all finite local PIRs with a prime characteristic. We start with a folk-law result:

Lemma 2.1 *If* $(R, \mathfrak{m}, \mathfrak{k})$ *is an artinian local ring such that* $\mathfrak{k}$ *is a finite field, then there exist a prime number* p *and positive integers* $r \leq s \leq t$ *such that* $char(R) = p^r$ *and* $|R| = p^t$*, where* s *is the nilpotency index of* $\mathfrak{m}$.

Proof. Assume that $\mathfrak{m}$ has a nilpotency index s and consider the following sequence of nonzero finite dimensional $\mathfrak{k}$-vector spaces

$$\mathfrak{m}^{s-1}, \mathfrak{m}^{s-1}/\mathfrak{m}^{s-2}, \cdots, \mathfrak{m}/\mathfrak{m}^2.$$

If $char(\mathfrak{k}) = p$, then $|R/\mathfrak{m}| = p^m$ and $|\mathfrak{m}| = p^n$ for some natural numbers m and n. Clearly, $|R| = p^{m+n}$ and $n \geq m(s-1)$. Since $p^s = 0$, it follows that $char(R) = p^r$ for some $r \leq s$. ∎

Now assume that $(R, \mathfrak{m})$ is a finite local PIR, where $\mathrm{char}(R/\mathfrak{m}) = p$, $\mathfrak{m} = R\alpha$ and α has nilpotency index $s+1$. By Lemma 2.1, the characteristic of $(R, \mathfrak{m})$ is p^r for some r with $1 \leq r \leq s+1$. Then $\mathbb{Z}_{p^r}$ is a subring of R and $p \in R\alpha$. Let $\varphi : \mathbb{Z}_{p^r} \to R/\mathfrak{m}$ be the natural map and set $F = \mathrm{im}(\varphi)$. Then $F = \{\overline{0}, \overline{1}, \cdots, \overline{p-1}\} \cong \mathbb{Z}_p$, and F is the prime subfield of the residue field $R/\mathfrak{m}$. Assume further $R/\mathfrak{m} = F[\overline{\beta}]$ with $\beta \in R$. Then

$$R = \mathbb{Z}_{p^r}[\beta] + \mathbb{Z}_{p^r}[\beta]\alpha + \mathbb{Z}_{p^r}[\beta]\alpha^2 + \cdots + \mathbb{Z}_{p^r}[\beta]\alpha^s = \mathbb{Z}_{p^r}[\beta, \alpha]$$

and there is a surjective ring homomorphism

$$\pi : \mathbb{Z}_{p^r}[X, Y] \to R = \mathbb{Z}_{p^r}[\beta, \alpha], f(X, Y) \mapsto f(\beta, \alpha).$$

Hence each finite local PIR is a proper homomorphic image of the polynomial ring $\mathbb{Z}_{p^r}[X, Y]$, if $\mathrm{char}(R) = p^r$. The problem is to determine the kernel of π. In this section, we consider the question for the case $\mathrm{char}(R) = p$.

By [2, Theorem 9] or [3, Theorems 7.7, 7.8], a complete noetherian local ring $(R, \mathfrak{m})$ contains a coefficient field, if R has the same characteristic as its residue field $R/\mathfrak{m}$. It holds in particular for every finite local ring R with characteristic p. For the ring R, we can construct a coefficient field of R starting from any generator of the multiplicative cyclic group $(R/\mathfrak{m})^*$.

Theorem 2.2. *For a finite local ring $(R, \mathfrak{m})$ with characteristic p, assume that $|R/\mathfrak{m}| = p^m$. Then there exist an element $\beta \in R$ and a subring $A = \mathbb{Z}_p[\beta]$ of R such that $\beta^{p^{mt}-1} = 1$ for some positive integer t and $R/\mathfrak{m} = \overline{A}$. In the case, A is a coefficient field of R.*

Proof. Denote $R/\mathfrak{m}$ by F and set $k = p^m$. Then F^* is a cyclic multiplicative group of order $k - 1$. Now let α be a generator of the group F^*. Then $\alpha^k = \alpha$, $\alpha^{k^t} = \alpha$ and hence $\alpha^{k^t-1} = 1$ holds for any positive integer t. Now let s be the nilpotency index of $\mathfrak{m}$ and choose t such that $k^t \geq s$. Choose $\beta_1 \in R$ such that $\overline{\beta_1} = \alpha$. Assume that $\beta_1^{k^t-1} = 1 + x$ for some $x \in \mathfrak{m}$. Clearly

$$(1 + x)^{k^t-1} = 1 + \sum_{r=1}^{s-1} C_{k^t-1}^r x^r.$$

Since $C_{k^t-1}^r \cdot x^r = C_{k^t}^r \cdot x^r + (-1)C_{k^t-1}^{r-1} \cdot x^r = (-1)C_{k^t-1}^{r-1} \cdot x^r$, it follows by induction that $C_{k^t-1}^r \cdot x^r = (-1)^r x^r$. Then $(1+x)^{k^t-1} = 1 + \sum_{r=1}^{s-1}(-1)^r x^r$. In this case, set $\beta = \beta_1(1+x)$. Then $\overline{\beta} = \alpha$ in $R/\mathfrak{m}$ and

$$\beta^{k^t-1} = \beta_1^{k^t-1}(1 + x)^{k^t-1} = (1 + x)(1 + \sum_{r=1}^{s-1}(-1)^r x^r) = 1.$$

Set $A = \{0, 1, \beta, \beta^2, \cdots, \beta^{k^t-2}\}$. Clearly, $R/\mathfrak{m} = \overline{A}$. We will show that A is a subring of R and thus a field under the operations in R. First, note that for any $\delta \in R$, there exist $\gamma \in A$ and $y \in \mathfrak{m}$ such that $\delta = \gamma + y$. Consider the equation $X^{k^t} = X$. Clearly, every element in A is a root of the equation. If $\delta \notin A$, then there exist $\gamma \in A$ and nonzero $y \in \mathfrak{m}$ such that $\delta = \gamma + y$, and so $\delta^{k^t} = (\gamma + y)^{k^t} = \gamma^{k^t} + y^{k^t} = \gamma = \delta - y \neq \delta$. This implies that δ is not a root of the equation, and so for any $\delta \in R$, $\delta \in A$ if and only if δ is a root of the equation $X^{k^t}=X$. Given any $\delta_1,\delta_2\in A$, then $(\delta_1-\delta_2)^{k^t}=\delta_1^{k^t}+(-1)^{k^t}\delta_2^{k^t}=\delta_1-\delta_2$, implying $\delta_1-\delta_2\in A$. Thus A is a subring of R and $A=\mathbb{Z}_p[\beta]$. Clearly, A is also a field under the operation and thus $A=\{0,1,\beta,\beta^2,\cdots,\beta^{k-2}\}$. ■

We now characterize finite local PIRs with characteristic p. The following corollary also follows from [5, Theorem 3.1], which is a direct consequence of Cohen's Theorem 9 ([2]).

Corollary 2.3. *Let $(R,\mathfrak{m})$ be a finite local ring with $\operatorname{char}(R)=p$. Then R is a PIR, $\mathfrak{m}$ has nilpotency index s if and only if $R\cong F[X]/(X^s)$ for some finite field F, where X is an indeterminant over F. In particular, R has two (respectively, three) nontrivial ideals if and only if $R\cong F[X]/(X^s)$ for some finite field F, where $s=3$ ($s=4$, respectively).*

Proof. Clearly, R is a PIR ring if $R\cong F[X]/(X^s)$ for any finite or infinite field F.

Conversely, assume that R is a PIR and a local ring with characteristic p. Assume further $J(R)=R\alpha$, where $\alpha^s=0,\alpha^{s-1}\neq 0$. Let A be the subfield of R as in Theorem 2.2. Then $R/R\alpha=\overline{A}$ and it implies $A+R\alpha^2+\cdots+R\alpha^{s-1}=R$. Thus $R\alpha=A\alpha+A\alpha^2+\cdots+A\alpha^{s-1}$, and finally $R=A[\alpha]$. Since $\alpha^s=0,\alpha^{s-1}\neq 0$, it follows that $R\cong A[X]/(X^s)$. □

3 Finite local PIRs with characteristic p^r $(r\geq 2)$

Throughout this section, assume that $(R,\mathfrak{m})$ is a local ring with $\operatorname{Char}(R)=p^r$, where $r\geq 2$ and $\mathfrak{m}$ has nilpotency index $s+1$. Assume further $\mathfrak{m}=R\alpha$, $R/\mathfrak{m}=F[\overline{\beta}]$ for some $\beta\in R$, and $F\cong\mathbb{Z}_p$. Then $\alpha^{s+1}=0$ and $\alpha^s\neq 0$. We are going to determine the kernel of π, where

$$\pi:\mathbb{Z}_{p^r}[X,Y]\to R=\mathbb{Z}_{p^r}[\beta,\alpha], f(X,Y)\mapsto f(\beta,\alpha).$$

For any $f(X)=\sum_i a_iX^i$ in the polynomial ring $\mathbb{Z}_{p^r}[X]$, set $\overline{f}(X)=\sum_i\varphi(a_i)X^i$, where $\varphi:\mathbb{Z}_{p^r}\to R/\mathfrak{m}$ is the natural map. Let $g(X)$ be a monic polynomial of $\mathbb{Z}_{p^r}[X]$ such that $\overline{g}(X)\in F[X]$ is the minimal polynomial of $\overline{\beta}$ over F, where $\deg(g(X))=\deg(\overline{g}(X))$ and $F=\operatorname{im}(\varphi)$. For any monic polynomial $g_1(X)\in\mathbb{Z}_{p^r}[X]$, note that $\overline{g_1}(X)=\overline{g}(X)\in F[X]$ if and only if $g(X)-g_1(X)\in p\mathbb{Z}_{p^r}[X]$, which is denoted as

$$g(X)\equiv g_1(X)(\mod p).$$

Since clearly there exists a natural ring epimorphism from $\mathbb{Z}_{p^r}[\beta]$ to $F[\overline{\beta}]=R/\mathfrak{m}$ and the kernel is $p\mathbb{Z}_{p^r}[\beta]+g(\beta)\mathbb{Z}_{p^r}[\beta]$, it follows that

$$\mathfrak{m}=\mathbb{Z}_{p^r}[\beta]p+Z_{p^r}[\beta]g(\beta)+\sum_{t=1}^{s}\mathbb{Z}_{p^r}[\beta]\alpha^t,\quad U(R)=U(\mathbb{Z}_{p^r}[\beta])+\mathfrak{m}.\quad(\triangle)$$

These equalities will be used repeatedly in Section Four. Also, there exists a natural ring isomorphism

$$\psi : \mathbb{Z}_{p^r}[X]/(p, g(X)) \to F[\overline{\beta}] = R/\mathrm{m},\ \overline{f(X)} \mapsto \overline{f}(\beta).$$

In fact, consider $\sigma : \mathbb{Z}_{p^r}[X] \to F[\overline{\beta}]$, $f(X) \mapsto \overline{f}(\beta)$. For any $f(X)$, write

$$f(X) = g(X)h(X) + r(X) = g(X)h(X) + pr_1(X) + r_2(X),$$

where $\deg(r_i(X)) < \deg(g(X))$, and the coefficients of $r_2(X)$ is in $\{0, 1, \cdots, p-1\}$. If $\overline{f}(\beta) = 0$, then $\overline{r_2}(\beta) = 0$. Therefore $\overline{r_2}(X) = 0$ in $F[X]$ and hence $r_2(X) = 0$ holds in $\mathbb{Z}_{p^r}[X]$. This shows $\ker(\sigma) = (p, g(X))$ and hence ψ is an isomorphism .

The following two claims and an equality will be used repeatedly in what follows:

Claim (i): *For any $t(X) \in \mathbb{Z}_{p^r}[X]$, if $t(X)$ is not in the ideal generated by p and $g(X)$, then $t(\beta) \in U(R)$.*

Claim (ii): *$g(\beta) \in \mathfrak{m}$, hence $g(\beta)^{s+1} = 0$ and thus β is integral over $\mathbb{Z}_{p^r}$.*

$$\mathfrak{m} = \{0\} \cup U(R)\alpha \cup U(R)\alpha^2 \cup \cdots \cup U(R)\alpha^s. \quad (*)$$

We now prove the main result of this section.

Theorem 3.1. *Assume that p is a prime number and r is an integer with $r \geq 2$. Then a ring R is a finite local PIR with char$(R) = p^r$ and has exactly s nontrivial ideals, if and only if there exist integers $n \geq 1, m \geq 0$, a monic polynomial $g(X)$ in $\mathbb{Z}_{p^r}[X]$ and polynomials $u_i(X), v_j(X) \in \mathbb{Z}_{p^r}[X] \setminus (p, g(X))$ $(1 \leq i \leq n,\ 1 \leq j \leq m)$, where $\overline{g}(\overline{X})$ is irreducible in the polynomial ring $\mathbb{Z}_p[\overline{X}]$ (that is, in $\mathbb{Z}_{p^r}[X]/p\mathbb{Z}_{p^r}[X]$), deg$(g(X) =$ deg$(\,\overline{g}(\overline{X})\,)$, deg$(u_i(X)) <$ deg$(g(X))$ and deg$(v_j(X)) <$ deg$(g(X))$, such that R is isomorphic to a factor ring S/Q, where $S = \mathbb{Z}_{p^r}[X, Y]$ and*

$$Q = (pY^{s+1-t_1},\ Y^{s+1},\ p - \sum_{i=1}^{n} u_i(X)Y^{t_i},\ g(X) - \sum_{j=1}^{m} v_j(X)Y^{s_j})$$

with $1 \leq t_1 < t_2 < \cdots < t_n \leq s$, $1 \leq s_1 < s_2 < \cdots < s_m \leq s$ and $(r-1)t_1 \leq s < rt_1$.

Proof. $\Longleftarrow$: Assume that $R = S/Q$. Clearly, $p^r \cdot R = 0$, $Y^{s+1} \in Q$ and $Q \subseteq (p, g(X), Y)$. Also, there are natural ring isomorphisms

$$\mathbb{Z}_{p^r}[X, Y]/(p, g(X), Y) \cong \mathbb{Z}_{p^r}[X]/(p, g(X)) \cong \mathbb{Z}_p[\overline{X}]/(\overline{g}(\overline{X})),$$

where $Z_p[\overline{X}]/(\overline{g}(\overline{X}))$ is a finite field by assumption on $g(X)$. Set $\mathfrak{m} = (p, g(X), Y)/Q$. Clearly, $R/\mathfrak{m} \cong \mathbb{Z}_p[\overline{X}]/(\overline{g}(\overline{X}))$, $\mathfrak{m} = R\overline{Y}$, where $\overline{Y}^s \neq 0$, $\overline{Y}^{s+1} = 0$. It implies $p^{r-1} \cdot R \neq 0$ whence char$(R) = p^r$. Then $\mathfrak{m}$ is a nilpotent maximal ideal of R. Thus R has a unique maximal ideal $R\overline{Y}$, which is principal and has nilpotency index $s+1$, and thus by Lemma 2.1, R is a finite ring.

$\Longrightarrow$: Let $(R, \mathfrak{m}, \mathfrak{k})$ be a finite local PIR, where the nilpotency index is $s+1$ and $\mathfrak{m} = R\alpha$. Assume that $\mathrm{char}(R) = p^r$, where $r \geq 2$. Clearly, $p \in \mathfrak{m}$. By $(*)$ we can suppose that $p = u\alpha^t$, where $u \in U(R)$ and $1 \leq t \leq s$. Since $p^r \cdot 1_R = 0$ while $p^{r-1} \cdot 1_R \neq 0$, it follow that $(r-1)t \leq s < rt$. We only need to determine the kernel of π, where

$$\pi : \mathbb{Z}_{p^r}[X, Y] \to R = \mathbb{Z}_{p^r}[\beta, \alpha], f(X, Y) \mapsto f(\beta, \alpha).$$

Consider the aforementioned ring isomorphism

$$\psi : \mathbb{Z}_{p^r}[X]/(p, g(X)) \to F[\overline{\beta}] = R/\mathrm{m},\ \overline{f(X)} \mapsto \overline{f}(\overline{\beta}).$$

There exists a polynomial $u_1(X) \in \mathbb{Z}_{p^r}[X] \setminus (p, g(X))$ such that $\psi(\overline{u_1(X)}) = \overline{u}$. We can further assume that $deg(u_1(X)) < deg(g(X))$. Then $u_1(\beta) - u \in \mathfrak{m}$, that is, $u = u_1(\beta) + m_1$ for some $m_1 \in \mathfrak{m}$, hence $p = u_1(\beta)\alpha^t + m_1\alpha^t = u_1(\beta)\alpha^{t_1} + u_2\alpha^{t_2}$ for some $u_2 \in U(R)$ and some integer t_2, where $t_1 = t$ and $t_2 > t_1$. For the same reason, there exists a polynomial $u_2(X) \in \mathbb{Z}_{p^r}[X] \setminus (p, g(X))$ satisfying $deg(u_2(X)) < deg(g(X))$, such that $\psi(\overline{u_2(X)}) = \overline{u}_2$. If further $u_2\alpha^{t_2} \neq 0$, then there exist elements $u_3 \in U(R)$ and integer t_3 with $t_2 < t_3$ such that $p = u_1(\beta)\alpha^{t_1} + u_2(\beta)\alpha^{t_2} + u_3\alpha^{t_3}$. Continuing the process, we obtain that

$$p = u_1(\beta)\alpha^{t_1} + u_2(\beta)\alpha^{t_2} + \cdots + u_n(\beta)\alpha^{t_n}$$

where $u_i(x) \in \mathbb{Z}_{p^r}[X] \setminus (p, g(X))$ such that $deg(u_i(X)) < deg(g(X))$, $\forall i = 1, 2, \cdots, n$, and $1 \leq t = t_1 < t_2 < \cdots < t_n \leq s$.

Next we consider the possible value of $g(\beta)$. If $g(\beta) \neq 0$, by $(*)$ we can assume $g(\beta) = v\alpha^{s_1}$, where $1 \leq s_1 \leq s$ and $v \in U(R)$. Similarly, we have a polynomial $v_1(X) \in \mathbb{Z}_{p^r}[X] \setminus (p, g(X))$ such that $\psi(\overline{v_1(X)}) = \overline{v}$ and $deg(v_1(X)) < deg(g(X))$. Then $g(\beta) = v_1(\beta)\alpha^{s_1} + v_2\alpha^{s_2}$ with $v_2 \in U(R)$ and $s_1 < s_2$. If $v_2\alpha^{s_2} \neq 0$, then there exists a polynomial $v_2(X) \in \mathbb{Z}_{p^r}[X] \setminus (p, g(X))$ satisfying $deg(v_2(X)) < deg(g(X))$, such that $\psi(\overline{v_2(X)}) = \overline{v}_2$. Thus $g(\beta) = v_1(\beta)\alpha^{s_1} + v_2(\beta)\alpha^{s_2} + v_3\alpha^{s_3}$ where $s_2 < s_3$. Continuing the analysis, we obtain that

$$g(\beta) = v_1(\beta)\alpha^{s_1} + v_2(\beta)\alpha^{s_2} + \cdots + v_m(\beta)\alpha^{s_m}$$

where $v_j(X) \in \mathbb{Z}_{p^r}[X] \setminus (p, g(X))$ such that $deg(v_j(X)) < deg(g(X))$, $\forall j = 1, 2, \cdots, m$ and $1 \leq s_1 < s_2 < \cdots < s_m \leq s$.

Take

$$Q = (pY^{s+1-t_1},\ Y^{s+1},\ p - \sum_{i=1}^{n} u_i(X)Y^{t_i},\ g(X) - \sum_{j=1}^{m} v_j(X)Y^{s_j})$$

Notice that if $g(\beta) = 0$, we substitute $g(X) - \sum_{j=1}^{m} v_j(X)Y^{s_j}$ by $g(X)$, and in this case, $m = 0$.

Clearly, $Q \subseteq ker(\pi)$. Conversely, for any $f(X, Y) \in \mathbb{Z}_{p^r}[X, Y]$, write

$$f(X, Y) \equiv Y^s w_s(X) + Y^{s-1}w_{s-1}(X) + \cdots + Yw_1(X) + w_0(X) (\mod Q) \quad (**)$$

where $w_i(X) \in \mathbb{Z}_{p^r}[X]$ such that $deg(w_i(X)) < deg(g(X))$. Now assume that $f(\beta, \alpha) = 0$. Then $\alpha^s w_0(\beta) = 0$ and hence $w_0(\beta) \in \mathfrak{m}$, that is, $w_0(X) \in (p, g(X))$, which together with $deg(w_i(X)) < deg(g(X))$ yield that $w_0(X) = pw_0'(X)$. Thus we can substitute $w_0(X)$ by $pw_0(X)$ in $(**)$ and then we have

$$f(X,Y) \equiv Y^s w_s(X) + Y^{s-1} w_{s-1}(X) + \cdots + Y w_1(X) (\mod Q) \quad (***)$$

so that $\alpha^s w_1(\beta) = 0$, hence $w_1(x) = pw_1'(x)$ for some $w_1'(x) \in \mathbb{Z}_{p^r}[X]$ with $deg(w_1'(x)) < deg(g(x))$. Thus we can substitute $w_1(X)$ by $pw_1(X)$ in $(***)$ and get that

$$f(X,Y) \equiv Y^s w_s(X) + Y^{s-1} w_{s-1}(X) + \cdots + Y^2 w_2(X) (\mod Q).$$

Since s is finite, we can reach

$$f(X,Y) \equiv 0 (\mod Q).$$

after finite steps. Therefore, $f(X,Y) \in Q$. This shows that $ker(\pi) = Q$, as required. ■

We remark that by a similar discussion as above, we can get an alternative description for finite local PIRs R with char$(R) = p$, which is essentially the same with Corollary 2.3.

Theorem 3.2. *Let p be a prime number, and let $(R, \mathfrak{m})$ be a finite local ring with char$(R) = p$. Then R is a PIR with exactly s nontrivial ideals if and only if there exist integers $m \geq 0$, a monic irreducible polynomial $g(X)$ in $\mathbb{Z}_p[X]$ and polynomials $v_j(X) \in \mathbb{Z}_p[X] \setminus (g(X))$ $(1 \leq j \leq m)$, where $deg(v_j(X)) < deg(g(X))$, such that R is isomorphic to a factor ring S/Q, where $S = \mathbb{Z}_p[X,Y]$ and*

$$Q = (Y^{s+1},\, g(X) - \sum_{j=1}^{m} v_j(X) Y^{s_j})$$

with $1 \leq s_1 < s_2 < \cdots < s_m \leq s$.

4 Finite local rings with at most three nontrivial ideals

Note that in Theorem 3.1, the ideal Q is generated by four polynomials. In some cases, it can be generated by less polynomials. In this section, we study the structure of finite local rings with at most three nontrivial ideals. It is easy to see that the maximal ideal of such a ring R is principal and hence R is a finite local PIR in each case. We will show that each Q can be generated by at most three polynomials. The detailed verifications will be omitted in most cases.

Let p be any prime number. For any positive integer n, it is well-known that there exists in $\mathbb{Z}_p[X]$ an irreducible monic polynomial $g(X)$ of degree n. In the following theorems, for any $f(X) = X^t + \sum_{i=0}^{t-1} a_i X^i \in \mathbb{Z}_p[X]$, set $f_r(X) = X^t + \sum_{i=0}^{t-1} a_i X^i \in \mathbb{Z}_{p^r}[X]$, where $1 \leq r \leq 4$.

In [6], it is proved that a local ring R has exactly one nontrivial ideal if and only if either $R \cong F[X]/(X^2)$ for some field F, or $R \cong V/p^2V$, where V is a discrete valuation ring of *characteristic zero* and residue field of *characteristic* p, for some prime number p. For a finite local ring, use the discussions of Theorems 3.1 and 3.2 we can obtain the following result.

Theorem 4.1. *R is a finite local ring with exactly one nontrivial ideal and residue field of characteristic p, if and only if R is isomorphic to one of the of following homomorphic images of a polynomial ring $\mathbb{Z}_{p^r}[X, Y]$ $(1 \leq r \leq 2)$:*
(1) $\mathbb{Z}_p[X, Y]/(g(X),\, Y^2)$
(2) $\mathbb{Z}_{p^2}[X]/(g(X) + pw(X))$
where $g(X)$ is monic and irreducible in $\mathbb{Z}_p[X]$ and $deg(w(X)) < deg(g(X))$.

Theorem 4.2. *R is a finite local ring with exactly two nontrivial ideals and residue field of characteristic p, if and only if R is isomorphic to one of the following homomorphic images of a polynomial ring $\mathbb{Z}_{p^r}[X, Y]$ $(1 \leq r \leq 3)$:*
(1) $\mathbb{Z}_p[X, Y]/(g(X),\, Y^3)$
(2) $\mathbb{Z}_{p^2}[X, Y]/(g_2(X) + pw_2(X),\, p - u_2(X)Y^2,\, pY)$ *or*
$\mathbb{Z}_{p^2}[X, Y]/(g_2(X) - v_2(X)Y + pw_2(X),\, p - u_2(X)Y^2,\, pY)$
(3) $\mathbb{Z}_{p^3}[X]/(g_3(X) + pz_3(X) + p^2w_3(X))$
where $g(X)$ is a monic and irreducible polynomial of $\mathbb{Z}_p[X]$, $z(X), w(X), u(X)$ and $v(X)$ are polynomials of $\mathbb{Z}_p[X]$ such that $deg(x(X)) < deg(g(X)), \forall x(X) \in \{z(X), w(X)\}$, $u(X) \not\equiv 0(\mod g(X))$ and $v(X) \not\equiv 0(\mod g(X))$.

We give an outlined proof to the following:

Theorem 4.3. *R is a finite local ring with exactly three nontrivial ideals and residue field of characteristic p, if and only if R is isomorphic to one of the following homomorphic images of a polynomial ring $\mathbb{Z}_{p^r}[X, Y]$ $(1 \leq r \leq 4)$:*
(1) $\mathbb{Z}_p[X, Y]/(g(X),\, Y^4)$
(2) $\mathbb{Z}_{p^2}[X, Y]/(g_2(X) + pw_2(X),\, Y^3 - pu_2(X),\, pY)$ *or*
$\mathbb{Z}_{p^2}[X, Y]/(g_2(X) - v_2(X)Y^2 + pw_2(X),\, Y^3 - pu_2(X),\, pY)$ *or*
$\mathbb{Z}_{p^2}[X, Y]/(g_2(X) - v_2(X)Y + pw_2(X),\, Y^3 - pu_2(X),\, pY)$ *or*
$\mathbb{Z}_{p^2}[X, Y]/(g_2(X) - pYv_2(X),\, Y^2 - pu_2(X) - pYz_2(X))$ *or*
$\mathbb{Z}_{p^2}[X, Y]/(g_2(X) - pw_2(X) - pYy_2(X),\, Y^2 - pu_2(X) - pYz_2(X))$ *or*
$\mathbb{Z}_{p^2}[X, Y]/(g_2(X) - v_2(X)Y - pYy_2(X) - pw_2(X),\, Y^2 - pu_2(X) - pYz_2(X))$
(3) $\mathbb{Z}_{p^4}[X]/(g_4(X) + py_4(X) + p^2z_4(X) + p^3w_4(X))$

where $g(X)$ is a monic and irreducible polynomial of $\mathbb{Z}_p[X]$, $y(X), w(X), z(X), u(X)$ and $v(X)$ are polynomials of $\mathbb{Z}_p[X]$ such that

$$deg(x(X)) < deg(g(X)), \forall x(X) \in \{y(X), z(X), w(X)\},$$

$u(X) \not\equiv 0(\mod g(X))$ *and* $v(X) \not\equiv 0(\mod g(X))$.

Proof. $\Longleftarrow$: In case (1) and (2) (respectively, case (3)), $\overline{(p, g(X), Y)}$ (respectively, $\overline{(p, g(X))}$) is the principal maximal ideal of R, and it has nilpotency index 4. Thus each ring is finite local and has three nontrivial ideals.

$\Longrightarrow$: If $(R, R\alpha)$ is a finite local ring and α has nilpotency index four, then $\mathrm{char}(R) \in \{p^i \,|\, 1 \leq i \leq 4\}$. Assume $R/R\alpha = \mathbb{Z}_p[\overline{\beta}]$, then $R = \mathbb{Z}_{p^i}[\beta, \alpha]$. Let $g(X)$ be the minimal polynomial of $\overline{\beta}$ in $\mathbb{Z}_p[X]$. Then $g(\beta) \in R\alpha$, where

$$R\alpha = U(R)\alpha \cup U(R)\alpha^2 \cup U(R)\alpha^3 \cup \{0\}.$$

Consider the ring epimorphism

$$\pi : \mathbb{Z}_{p^i}[X, Y] \to R = \mathbb{Z}_{p^i}[\beta, \alpha],\ f(X, Y) \mapsto f(\beta, \alpha).$$

Case 1: $\mathrm{char}(R) = p$.

In this case, the result is stated in (1) and it follows from Corollary 2.3.

Case 2: $\mathrm{char}(R) = p^2$. In this case, either $p \in U(R)\alpha^2$ or $p \in U(R)\alpha^3$.

Subcase 2.1: $p \in U(R)\alpha^3$. Under the assumption, we have $p\mathfrak{m} = 0$ and thus there exists a polynomial $u(X) \in \mathbb{Z}_p[X] \setminus (g(X))$ such that $Y^3 - pu_2(X) \in \ker(\pi)$. If further $g_2(\beta) \in U(R)\alpha^3 \cup \{0\}$, then

$$\ker(\pi) = (g_2(X) + pw_2(X),\ Y^3 - pu_2(X),\ pY)$$

for some $w(X) \in \mathbb{Z}_p[X]$, where $\deg(w(X)) < \deg(g(X))$. If further $g_2(\beta) \in U(R)\alpha^2$, then

$$\ker(\pi) = (g_2(X) - v_2(X)Y^2 + pw_2(X),\ Y^3 - pu_2(X),\ pY),$$

where $v(X) \not\equiv 0(\mod g(X))$ in $\mathbb{Z}_p[X]$. If $g_2(\beta) \in U(R)\alpha$, then

$$\ker(\pi) = (g_2(X) - v_2(X)Y + pw_2(X),\ Y^3 - pu_2(X),\ pY).$$

Subcase 2.2: $p \in U(R)\alpha^2$. In this case, it follows by ($\triangle$) that

$$Y^2 - pu_2(X) - pYz_2(X) \in \ker(\pi)$$

for some $u(X) \in \mathbb{Z}_p[X] \setminus (g(X))$, where either $z(X) = 0$ or $z(X) \in \mathbb{Z}_p[X] \setminus (g(X))$.

If further $g_2(\beta) \in U(R)\alpha^3$, then by ($\triangle$) we have $g_2(X) - pYv_2(X) \in \ker(\pi)$ for some $v(X) \in \mathbb{Z}_p[X] \setminus (g(X))$, where $\deg(v(X)) < \deg(g(X))$. Set

$$Q_4 = (g_2(X) - pYv_2(X),\ Y^2 - pu_2(X) - pYz_2(X)).$$

Then clearly $Q_4 \subseteq \ker(\pi)$, $\{Y^4, pY^2, pg_2(X), Yg_2(X)\} \subseteq Q_4$ and $Y^3 \equiv t_*(X)(\mod Q_4)$. Conversely, $\forall f(X,Y) \in \ker(\pi)$,i.e., $f(\beta,\alpha)=0$, we have

$$f(X,Y) \equiv Y^3 t_*(X) + Y^2 t_*(X) + Y t_*(X) + t_*(X)(\mod Q_4)$$

$$\equiv Yt(X) + s(X)(\mod Q_4)$$

for some $t(X), s(X) \in \mathbb{Z}_{p^2}[X,Y]$, where $\deg(t(X)) < \deg(g(X))$ and $\deg(s(X)) < \deg(g(X))$. Then by Claim (i), $\alpha t(\beta) + s(\beta) = 0$ implies $t(X) \equiv 0(\mod p)$ and $s(X) \equiv 0(\mod p)$. Hence $f(X,Y) \equiv p \cdot (Y \cdot x(X) + y(X))(\mod Q_4)$, where the degree of both $x(X)$ and $y(X)$ are less than $\deg(g(X))$. It implies $Yy(X) \equiv 0(\mod Q_4)$, and therefore, $y(X) \equiv 0(\mod p)$. Then $f(X,Y) \equiv (pY) \cdot x(X)(\mod Q_4)$, and thus $x(X) \equiv 0(\mod p)$ by Claim (i). Finally, $f(X,Y) \in Q_4$, and it implies $Q_4 = \ker(\pi)$ in the case.

If further $g_2(\beta) \in U(R)\alpha^2 \cup \{0\}$, then it follows from $(\triangle)$ that

$$\ker(\pi) = (g_2(X) - pw_2(X) - pYy(X),\, Y^2 - pu_2(X) - pYz_2(X))$$

holds for some $y(X), z(X), w(X) \in \mathbb{Z}_p[X]$ with

$$\deg(x(X)) < \deg(g(X)),\ \forall x(X) \in \{y(X), z(X), w(X)\}.$$

If further $g_2(\beta) \in U(R)\alpha$, then it follows from $(\triangle)$ that

$$\ker(\pi) = (g_2(X) - v_2(X)Y - pYy_2(X) - pw_2(X),\, Y^2 - pu_2(X) - pYz_2(X)),$$

where $u(X), v(X), w(X), y(X)$ and $z(X)$ has the above mentioned properties.

Case 3: $\mathrm{char}(R) = p^3$.

The situation can not occur since $p \in U(R)\alpha \cup U(R)\alpha^2 \cup U(R)\alpha^3$.

Case 4: $\mathrm{char}(R) = p^4$.

In this case, $p \in U(R)\alpha$, thus α is nothing new but a unital multiple of p in $\mathbb{Z}_{p^4}[X]$. In this case, it is easy to obtain (3) by $(\triangle)$. ■

In the following Propositions 4.4 and 4.5, assume that $g_i(X)$ and $u_j(X)$ are monic polynomials in $\mathbb{Z}_{p^2}[X]$ such that $\deg(u_r(X)) < \deg(g_1(X))$ $(r = 1,3)$, $\deg(u_s(X)) < \deg(g_2(X))$ $(s = 2,4)$, $g_i(\overline{X})$ are irreducible in $\mathbb{Z}_p[\overline{X}]$ (i.e., $\mathbb{Z}_{p^2}[X,Y]/(p, g_i(X))$) and $u_j(X) \not\equiv 0(\mod p)$. We can assume further that the coefficients of $u_j(X)$ and $g_i(X)$ are in $\{0,1,\cdots,p-1\}$ $(1 \le i \le 2,\ 1 \le j \le 4)$. Then either $g_1(X) = g_2(X)$ or there exist $u(X), v(X)$ such that

$$g_1(X)u(X) + g_2(X)v(X) \equiv 1(\mod p).$$

In Proposition 4.4, set for $i = 1,2$

$$R_i = \mathbb{Z}_{p^2}[X,\, Y]/(g_i(X),\, Y^2 - p \cdot u_i(X),\, pY),$$

$$\mathfrak{m}_i = (p,\, g_i(X),\, Y)/(g_i(X),\, Y^2 - p\cdot u_i(X),\, pY)$$

and $K_i = \mathbb{Z}_{p^2}[X]/(p,\, g_i(X)) = \mathbb{Z}_p[\overline{X}]/(g_i(\overline{X}))$. Let $\pi_i : R_i \to K_i$ be the natural epimorphism. Note that $\mathfrak{m}_i$ is the maximal ideal of R_i for each i. Note also that $K_1 \cong K_2$ if and only if $\deg(g_1(X)) = \deg(g_2(X))$.

Proposition 4.4. (1) *If R_1 is isomorphic to R_2, then there exist a field isomorphism $\tau : K_1 \to K_2$, a polynomial $v_2(X)$ in $\mathbb{Z}_{p^2}[X]\backslash(p, g_2(X))$ such that $\overline{u_2(X)} = \overline{v_2(X)}^2 \cdot \tau(\,\overline{u_1(X)}\,)\,)$ holds in K_2.*

(2) *Conversely, if $g_1(X) = g_2(X)$ and there exist $v_i(X)$ in $\mathbb{Z}_{p^2}[X]$ such that $\overline{u_2(X)} = \overline{v_2(X)}^2 \cdot \overline{u_1(X)}$ holds in K_2 and $\overline{u_1(X)} = \overline{v_1(X)}^2 \cdot \overline{u_2(X)}$ holds in K_1 respectively, then R_1 is isomorphic to R_2.*

Proof. (1) Assume that $\sigma : R_1 \to R_2$ is a ring isomorphism. Then $\sigma(\mathfrak{m}_1) = \mathfrak{m}_2$, and hence σ induces a natural field isomorphism τ from K_1 to K_2. Clearly $\mathfrak{m}_i = R_i\overline{Y}$, and then $\sigma(R_1\overline{Y}) = R_2\overline{Y}$. Let $\overline{v(X,Y)}\cdot\sigma(\overline{Y}) = \overline{Y}$ for some $v(X,Y) \in \mathbb{Z}_{p^2}[X,Y]$. Note that $\overline{Y}^2 = p\cdot\overline{u_i(X)}$ in R_i $(i = 1, 2)$. Thus in R_2

$$p\cdot\overline{u_2(X)} = \overline{v(X,Y)}^2\sigma(\overline{Y}^2) = p\cdot\overline{v(X,Y)}^2\sigma(\overline{u_1(X)}).$$

Note that $p\cdot\overline{Y} = 0$ in R_2, thus assume

$$p\cdot\sigma(\,\overline{u_1(X)}\,) = p\cdot\overline{w(X)}, \quad p\cdot\overline{v(X,Y)} = p\cdot\overline{v_2(X)}$$

for some $w(X), v_2(X) \in \mathbb{Z}_{p^2}[X]$. Then $p\cdot[u_2(\,\overline{X}\,) - v_2(\,\overline{X}\,)^2\cdot w(\,\overline{X}\,)] = 0$ holds in the *cc*-local ring $(R_2,\, R_2\overline{Y})$, where $R_2 = \mathbb{Z}_{p^2}[\overline{X}, \overline{Y}]$. By *Claim* (*i*), we must have $u_2(X) - v_2(X)^2\cdot w(X) \in (p,\, g_2(X))$. Consider the following commutative diagram in which the isomorphism τ is induced by σ.

$$\begin{tikzcd}
R_1 \arrow[r, "\sigma"] \arrow[d, "\pi_1"'] & R_2 \arrow[d, "\pi_2"] \\
K_1 \arrow[r, "\tau"'] & K_2
\end{tikzcd}$$

We get $\overline{u_2(X)} = \overline{v_2(X)}^2 \cdot \tau(\,\overline{u_1(X)}\,)\,)$ in K_2. Since $\overline{u_2(X)} \in U(R_2)$, both $\overline{v_2(X)}$ and $\tau(\,\overline{u_1(X)})$ are invertible in R_2. Note that $\overline{u_1(X)} = \overline{v_1(X)}^2 \cdot \tau^{-1}(\,\overline{u_2(X)}\,)\,)$ holds in K_1 for some $v_1(X) \in \mathbb{Z}_{p^2}[X]$.

(2) By assumption, we can assume $\overline{w_1(X)}^2 \cdot \overline{u_2(X)} = \overline{u_1(X)}$ in K_2. Define a ring homomorphism $\varphi_1 : \mathbb{Z}_{p^2}[X,Y] \to R_2$ by

$$1 \mapsto 1,\ \varphi(X) = X,\ \varphi(Y) = \overline{w_1(X)Y}.$$

Since $p\cdot\overline{(p, g_2(X))} = 0$ in R_2,

$$\varphi_1(Y^2 - pu_1(X)) = \overline{w_1(X)Y}^2 - p\overline{u_1(X)} = \overline{w_1(X)}^2\cdot(\overline{Y}^2 - p\overline{u_2(X)}) = 0$$

holds in R_2. Then $(g_1(X),\, Y^2 - p \cdot u_1(X),\, pY) \subseteq \ker(\varphi_1)$, and φ_1 induces a ring homomorphism $\overline{\varphi_1} : R_1 \to R_2$ such that

$$\overline{\varphi_1}(\overline{X}) = \overline{X}, \quad \overline{\varphi_1}(\overline{Y}) = \overline{w_1(X)} \cdot \overline{Y}.$$

Under the assumption, there exists another ring homomorphism $\overline{\varphi_2} : R_2 \to R_1$ such that $\overline{\varphi_2}(\overline{X}) = \overline{X}$, $\overline{\varphi_1}(\overline{Y}) = \overline{w_2(X)} \cdot \overline{Y}$. Then $(\overline{\varphi_2} \cdot \overline{\varphi_1})(\overline{Y}) = (\,\overline{w_1(X)} \cdot \overline{w_2(X)}\,) \cdot \overline{Y}$, and therefore $\overline{\varphi_2} \cdot \overline{\varphi_1}$ is a ring isomorphism. ■

In the following Proposition 4.5, Set $R_1 = S/\mathfrak{I}_1$, $R_2 = S/\mathfrak{I}_2$, where $S = \mathbb{Z}_{p^2}[X,\, Y]$,

$$\mathfrak{I}_1 = (g_1(X) - u_3(X)Y,\, Y^2 - p \cdot u_1(X), pY)$$

$$\mathfrak{I}_2 = (g_2(X) - u_4(X)Y,\, Y^2 - p \cdot u_2(X),\, pY).$$

Set $K_i = \mathbb{Z}_{p^2}[X]/(p,\, g_i(X)) \cong \mathbb{Z}_p[X]/(\overline{g}_i(X))$. Let $\pi_i : R_i \to K_i$ be the natural epimorphism. Note that $K_1 \cong K_2$ if and only if $\deg(g_1(X)) = \deg(g_2(X))$. By the proof of Proposition 3.2, we have

Proposition 4.5. (1) *If R_1 is isomorphic to R_2, then there exist a field isomorphism $\tau : K_1 \to K_2$, polynomials $v_2(X)$, $u_5(X)$ in $\mathbb{Z}_{p^2}[X] \setminus (p, g(X))$ such that $\sigma(\,\overline{g_1(X)}\,) = \overline{u_5(X)} \cdot \overline{g_2(X)}$ holds in R_2 and*

$$\overline{u_2(X)} = \overline{v_2(X)}^2 \cdot \tau(\,\overline{u_1(X)})$$

hold in K_2.

(2) *Conversely, if $g_1(X) = g_2(X)$ and there exist $v_i(X)$ in $\mathbb{Z}_{p^2}[X]$ such that $u_2(X) \equiv v_2(X)^2 \cdot u_1(X)(\mod (p, g_2(X)^2))$, $u_4(X) \equiv v_2(X)u_3(X)(\mod (p, g_2(X)^2))$ and $u_1(X) = v_1(X) \cdot u_2(X)(\mod (p, g_1(X)^2)), u_3(X) \equiv v_1(X)u_4(X)(\mod (p, g_1(X)^2))$ respectively, then R_1 is isomorphic to R_2.*

References

[1] M.F. Atiyah and I.G. MacDonald. *Introduction to Commutative Algebra*, **Addison-Wesley**, Reading, MA, 1969.

[2] I.S. Cohen. On the structure and ideal theory of complete local rings. **Trans. Amer. Math. Soc. 59** (1946), 54 – 106.

[3] D. Eisenbud. *Commutative Algebra with a View Toward Algebraic Geometry.* **Springer Science + Business Media, Inc.**, 2004.

[4] D.C. Lu and T.S. Wu. The zero-divisor graphs which are uniquely determined by neighborhoods. *Comm. in Algebra*, **35**(2007), 3855 – 3864.

[5] K.R. McLean. Commutative artinian principal ideal rings. **Proc. London Math. Soc. 26** (1973), 249 – 272.

[6] J. Reineke. Commutative rings in which every proper ideal is maximal. **Fund. Math 97** (1977), 229-231.

[7] T.S. Wu and D.C. Lu. The structure of a class of Z-local rings. **Science in China Series A: Mathematics 49:10**(2006), 1297 – 1302.

[8] T.S. Wu, D.C. Lu and Y.L. Li. On zero divisors and prime elements of po-semirings. **Preprint**.